

---

# **PyGEDM**

**Jul 19, 2021**



---

## API Index:

---

<b>1</b>	<b>PyGEDM</b>	<b>3</b>
1.1	Web app . . . . .	3
1.2	Usage . . . . .	3
1.3	Installation . . . . .	4
1.4	References . . . . .	4
1.5	YMW16 C LICENSE . . . . .	5
	<b>Python Module Index</b>	<b>13</b>
	<b>Index</b>	<b>15</b>







# CHAPTER 1

## PyGEDM

*Python bindings for the YMW16, NE2001 and YT2020 electron density models*

This package is a Python interface to the YMW16 and NE2001 electron density models, and YT2020 halo model. The Yao, Manchester and Wang (2017, *Astrophys. J.*, 835, 29; arXiv:1610.09448) YMW16 electron density model, is written in C++, and the Cordes and Lazio (2001, arXiv:0207156) NE2001 model is written in FORTRAN. This package, PyGEDM, wraps these two codes using `pybind11` to make them usable from Python. Here, we have converted NE2001 to C++ using `f2c`.

### 1.1 Web app

We provide a web app at <https://apps.datacentral.org.au/pygedm/>

The pygedm web app is kindly hosted by [Data Central](#).

### 1.2 Usage

Some usage examples can be found in the `examples` directory.

```
import pygedm

# calculate DM at a given distance
DM, tau_sc = pygedm.dist_to_dm(204.0, -6.5, 200, method='ne2001')
DM, tau_sc = pygedm.dist_to_dm(204.0, -6.5, 200, method='ymw16')

# calculate distance for a given sky position and DM
dist, tau_sc = pygedm.dm_to_dist(123.4, 4.0, 200)

# calculate N_e density at xyz galactocentric coordinates
ne = pygedm.calculate_electron_density_xyz(1.0, 2.0, 3.0)
```

(continues on next page)

(continued from previous page)

```
# calculate N_e density at Galactic lat/long/distance coords
ne = pygedm.calculate_electron_density_lbr(204.0, -6.5, 3000.0)

# Calculate halo DM contribution
dm_halo = pygedm.calculate_halo_dm(gl=0, gb=30)
```

The methods return astropy [Quantities](#), which have units attached, and can accept astropy [Angles](#) and [Quantities](#) as arguments:

```
import pygedm
import astropy.units as u
import astropy.coordinates as c
DM = u.Quantity(10.0, unit='pc cm^-3')
ra, dec = c.Angle(23.0, unit='hourangle'), c.Angle('-43:00:02', unit='degree')
sky_coords = c.SkyCoord(ra, dec, frame='icrs')
dist, tau_sc = pygedm.dm_to_dist(sky_coords.galactic.l, sky_coords.galactic.b, DM)

print(dist.to('lyr'))
>> 3362.16343117 lyr
print(tau_sc.to('ns'))
>> 7.758686138 ns
```

## 1.3 Installation

Requires pybind11, astropy, numpy, scipy, a newish C compiler with C++11 support (Ubuntu 16.04+ default gcc will work), plus f2c.

Pre-compiled wheels are generated for linux, meaning you can skip the f2c install step. Alternatively you can install via apt-get f2c in Ubuntu, or via conda install -c conda-forge f2c if you use conda.

For MacOS, you are best off using conda and getting f2c via conda install -c conda-forge f2c. Windows is not currently supported.

You should be able to install with:

or

to install the latest version from github. Alternatively, download this repository and install via

To run unit tests, run python setup.py test. Note that these tests only check the Python bindings, not the underlying C/FORTRAN source code (which is not supplied with unit tests).

## 1.4 References

If using PyGEDM in a journal article, please remember to cite the underlying electron density models:

Cordes, J. M., & Lazio, T. J. W. (2002), *NE2001.I. A New Model for the Galactic Distribution of Free Electrons and its Fluctuations*, arXiv e-prints, astro-ph/0207156.

Cordes, J. M., & Lazio, T. J. W. (2003), *NE2001. II. Using Radio Propagation Data to Construct a Model for the Galactic Distribution of Free Electrons*, arXiv e-prints, astro-ph/0301598.

Yao, J. M., Manchester, R. N., & Wang, N. (2017), *A New Electron-density Model for Estimation of Pulsar and FRB Distances*, The Astrophysical Journal, Volume 888, Issue 2, id.105, Column 835, id.29

Yamasaki, S., & Totani, T. (2020), *The Galactic Halo Contribution to the Dispersion Measure of Extragalactic Fast Radio Bursts*, The Astrophysical Journal, Volume 888, Issue 2, id.105

These are available in bibtex format in [references.bib](#), and also as an [ADS library](#).

### 1.4.1 YMW16 C README

YMW16 is a model for the distribution of free electrons in the Galaxy, the Magellanic Clouds and the inter-galactic medium, that can be used to estimate distances for real or simulated pulsars and fast radio bursts (FRBs) based on their position and dispersion measure.

The Galactic model is based on 189 pulsars that have independently determined distances as well as dispersion measures, whereas simpler models are used for the electron density in the MC and the IGM. It is estimated that the 95% of predicted Galactic pulsar distances will have a relative error of less than a factor of 0.9. Pulse broadening due to scattering in the Galactic interstellar medium, the Magellanic Clouds, the intergalactic medium and FRB host galaxies is estimated.

As well as the ymw16 dm-distance program, we also provide a program, ymw16\_ne, which gives the electron density at any point in the Galaxy or Magellanic Clouds.

A paper (Yao, Manchester and Wang, 2017, *Astrophys. J.*, 835, 29; [arXiv:1610.09448](#)) describes the model and compares its predictions with those of earlier Galactic electron density models. YMW16 is the first electron-density model to estimate extragalactic pulsar distances and FRB distances.

To make a command-line executable version of the program, download and unpack the latest version of the program. Then run “make\_ymw16” to create the executable. The environment variable YMW16\_DIR may be set up to point at a directory containing ymw16par.txt and spiral.txt. Access to these files is needed at runtime.

Websites allowing interactive access to the YMW16 distance model and download of the latest program version are available at:

- <http://www.xao.ac.cn/ymw16/>,
- <http://www.atnf.csiro.au/research/pulsar/ymw16/> and
- <https://bitbucket.org/psrsoft/ymw16/>.

Please report any issues or bugs at [https://bitbucket.org/psrsoft/ymw16/issues/new/](https://bitbucket.org/psrsoft/ymw16/issues/new) or directly to the authors. Please provide an example illustrating the problem.

## 1.5 YMW16 C LICENSE

### 1.5.1 NE2001 README

07 July 2002 To compile and execute the code, see [code.pdf](#).

### pygedm.pygedm

Python API to YMW16, NE2001, and YT2020 Galactic electron density models

### References

[1] Cordes, J. M., & Lazio, T. J. W. (2002), *NE2001.I. A New Model for the Galactic Distribution of Free Electrons and its Fluctuations*, arXiv e-prints, astro-ph/0207156.

[2] Cordes, J. M., & Lazio, T. J. W. (2003), *NE2001. II. Using Radio Propagation Data to Construct a Model for the Galactic Distribution of Free Electrons*, arXiv e-prints, astro-ph/0301598.

[3] Yao, J. M., Manchester, R. N., & Wang, N. (2017), *A New Electron-density Model for Estimation of Pulsar and FRB Distances*, ApJ, 835, 29.

[4] Yamasaki S, Totani T (2020), *The Galactic Halo Contribution to the Dispersion Measure of Extragalactic Fast Radio Bursts* The Astrophysical Journal, Volume 888, Issue 2, id.105

`pygedm.pygedm.calculate_electron_density_lbr(gl, gb, dist, method='ymw16')`

Calculate electron density at a point with Galactic coords (ga, gl) at given distance

### Parameters

- `gl` (*float, Angle, or Quantity*) – Galactic longitude in degrees (or astropy Angle)
- `gb` (*float, Angle, or Quantity*) – Galactic latitude in degrees (or astropy Angle)
- `dist` (*float or Quantity*) – Distance in pc

**Returns** electron density in cm<sup>-3</sup>

**Return type** N\_e (astropy.Quantity)

`pygedm.pygedm.calculate_electron_density_xyz(x, y, z, method='ymw16')`

Calculate electron density at a point with galactocentric coords (x, y, z)

### Parameters

- `x` (*float or Quantity*) – galactocentric X coordinate in pc
- `y` (*float or Quantity*) – galactocentric Y coordinate in pc
- `z` (*float or Quantity*) – galactocentric Z coordinate in pc

**Returns** electron density in cm<sup>-3</sup>

**Return type** N\_e (astropy.quantity)

`pygedm.pygedm.calculate_halo_dm(gl, gb, method='yt2020', component='both')`

Compute halo DM

### Parameters

- `gl` (*float, Angle, or Quantity*) – Galactic longitude in degrees (or astropy Angle)
- `gb` (*float, Angle, or Quantity*) – Galactic latitude in degrees (or astropy Angle)
- `method` (*str*) – one of ‘yt2020’ (only YT2020 supported currently)
- `component` (*str*) – Compute ‘spherical’ component of halo, ‘disk’, or ‘both’ components.

**Returns** Dispersion measure in (pc/cm<sup>3</sup>)

**Return type** DM (float)

`pygedm.pygedm.convert_lbr_to_xyz(gl, gb, dist, method='ymw16')`

Convert Galactic (l,b,r) coords to Galactocentric (x,y,z) coords

### Parameters

- `gl` (*float, Angle, or Quantity*) – Galactic longitude in degrees (or astropy Angle)
- `gb` (*float, Angle, or Quantity*) – Galactic latitude in degrees (or astropy Angle)

- **dist** (*float or Quantity*) – Distance in pc
- **method** (*str*) – one of ‘ymw16’, ‘ne2001’, or ‘astropy’

**Returns** Galactocentric X, Y, Z coordinates

**Return type** xyz (tuple)

## Notes

For transform, the Sun is located at (x=0, y=R\_sun, z=z\_sun) YMW16 uses R\_sun of 8300 pc and z\_sun of 6.0 pc NE2001 uses R\_sun of 8500 pc and z\_sun of 0.0 pc Both of these do a basic spherical to cartesian conversion.

astropy does a much more complicated conversion, see <https://astropy.readthedocs.io/en/latest/coordinates/galactocentric.html> This is the ‘proper’ coordinate system, but note that it is NOT COMPATIBLE WITH NE2001 OR YMW16! (!SEE EXAMPLE OUTPUT BELOW!)

Example output:

```
pygedm.convert_lbr_to_xyz(0, 0, 0, method='ymw16')
(<Quantity 0. pc>, <Quantity 8300. pc>, <Quantity 6. pc>)

pygedm.convert_lbr_to_xyz(0, 0, 0, method='ne2001')
(<Quantity 0. pc>, <Quantity 8500. pc>, <Quantity 0. pc>)

pygedm.convert_lbr_to_xyz(0, 0, 0, method='astropy')
(<Quantity -8499.95711754 pc>, <Quantity 0. pc>, <Quantity 27. pc>)
```

pygedm.pygedm.**dist\_to\_dm**(*gl, gb, dist, mode='gal', method='ymw16', nu=1.0*)

Convert a distance to a DM

### Parameters

- **gl** (*float in deg or astropy.Angle*) – galactic longitude
- **gb** (*float in deg or astropy.Angle*) – galactic latitude
- **dist** (*float or astropy.Quantity*) – distance to source (pc) or if in mode IGM use (Mpc)
- **method** (*str*) – choose electron density model, either ‘ymw16’ or ‘ne2001’
- **mode** (*str*) – Gal, MC, or IGM (for YMW16 only)
- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)

**Returns** Dispersion measure (pc / cm3), scattering timescale at 1 GHz (s)

**Return type** dm (astropy.Quantity), tau\_sc (astropy.Quantity)

pygedm.pygedm.**dm\_to\_dist**(*gl, gb, dm, dm\_host=0, mode='gal', method='ymw16', nu=1.0*)

Convert a DM to a distance

### Parameters

- **gl** (*float in deg or astropy.Angle*) – galactic longitude
- **gb** (*float in deg or astropy.Angle*) – galactic latitude
- **dm** (*float in pc/cm3 or astropy.Quantity*) – dispersion measure (pc cm^-3)
- **method** (*str*) – choose electron density model, either ‘ymw16’ or ‘ne2001’
- **mode** (*str*) – Gal, MC, or IGM (for YMW16 only)

- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)

**Returns** Distance (pc), scattering timescale at 1 GHz (s)

**Return type** dist (astropy.Quantity), tau\_sc (astropy.Quantity)

`pygedm.pygedm.generate_healpix_dm_map(dist=1, nside=64, method='ymw16')`

Generate an all-sky healpix map for a given distance.

### Parameters

- **dist** (*float or Quantity*) – Distance to integrate EDM out to. 30 kpc will go to edge
- **nside** (*int*) – The NSIDE parameter for the healpix map (power of 2, larger => higher resolution)
- **method** (*str*) – one of ‘ymw16’, ‘ne2001’, ‘yt2020’ or ‘yt2020\_analytic’

### Notes

This method takes a fair amount of time to run – tens of seconds for NSIDE=32. YT2020 method is even slower, consider using yt2020\_analytic

**Returns** Healpix map as a numpy array (1D), which can be viewed using the healpy.mollview() method

**Return type** hmap (np.array)

## pygedm.ne2001\_wrapper

Python wrapper for NE2001 model code.

### References

[1] Cordes, J. M., & Lazio, T. J. W. (2002), *NE2001.I. A New Model for the Galactic Distribution of Free Electrons and its Fluctuations*, arXiv e-prints, astro-ph/0207156.

[2] Cordes, J. M., & Lazio, T. J. W. (2003), *NE2001. II. Using Radio Propagation Data to Construct a Model for the Galactic Distribution of Free Electrons*, arXiv e-prints, astro-ph/0301598.

`pygedm.ne2001_wrapper.TAUISS(d, sm, nu)`

Convert a scattering measure (SM) to scattering timescale at given frequency.

Direct port from FORTRAN code scattering98.f

### Parameters

- **d** (*float*) – Distance in kpc
- **sm** (*float*) – Scattering measure (kpc m<sup>-20/3</sup>)
- **nu** (*float*) – Radio frequency in GHz

**Returns** pulse broadening time (ms)

**Return type** tauiss (float)

Fortran equiv:

```

REAL FUNCTION TAUSS(d, sm, nu)
c
c calculates the pulse broadening time in ms
c from distance, scattering measure, and radio frequency
c
c input:      d = pulsar distance      (kpc)
c                  sm = scattering measure   (kpc m^{-20/3})
c                  nu = radio frequency       (GHz)
c output: tausis = pulse broadening time (ms)
c
c      implicit none
real d, sm, nu
tauss = 1000. * (sm / 292.)**1.2 * d * nu**(-4.4)
end

```

`pygedm.ne2001_wrapper.calculate_electron_density_xyz(x, y, z)`

Compute electron density at Galactocentric X, Y, Z coordinates

x,y,z are Galactocentric Cartesian coordinates, measured in kpc (NOT pc!) with the axes parallel to (l, b) = (90, 0), (180, 0), and (0, 90) degrees

#### Parameters

- **x** (*float*) – Galactocentric coordinates in kpc
- **y** (*float*) – Galactocentric coordinates in kpc
- **z** (*float*) – Galactocentric coordinates in kpc

**Returns** Electron density in cm-3

**Return type** ne\_out (astropy.Quantity)

`pygedm.ne2001_wrapper.dist_to_dm(l, b, dist, nu=1.0, full_output=False)`

Convert distance to DM and compute scattering timescale

#### Parameters

- **l** (*float*) – galactic longitude in degrees
- **b** (*float*) – galactic latitude in degrees
- **dist** (*float*) – Distance in kpc
- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)
- **full\_output** (*bool*) – Return full raw output (dict) from NE2001 if set to True

**Returns** Dispersion measure (pc / cm3), scattering timescale at 1 GHz (s)

**Return type** dm (astropy.Quantity), tau\_sc (astropy.Quantity)

`pygedm.ne2001_wrapper.dm_to_dist(l, b, dm, nu=1.0, full_output=False)`

Convert DM to distance and compute scattering timescale

#### Parameters

- **l** (*float*) – galactic longitude in degrees
- **b** (*float*) – galactic latitude in degrees
- **dm** (*float*) – Dispersion measure
- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)
- **full\_output** (*bool*) – Return full raw output (dict) from NE2001 if set to True

**Returns** Distance (pc), scattering timescale at 1 GHz (s)

**Return type** dist (astropy.Quantity), tau\_sc (astropy.Quantity)

```
pygedm.ne2001_wrapper.run_from_pkgdir(f)
```

Decorator function to chdir() into package directory when running

NE2001 code doesn't know the relative path to its data files. This wraps the function call, changing into the right directory first, calling it, then changing back to original directory.

## pygedm.ymw16\_wrapper

Python/C++ port of YMW16 C code

### References

[1] Yao, J. M., Manchester, R. N., & Wang, N. (2017), *A New Electron-density Model for Estimation of Pulsar and FRB Distances*, ApJ, 835, 29.

```
pygedm.ymw16_wrapper.calculate_electron_density_lbr(gl, gb, dist)
```

**Calculate electron density at a point with Galactic coords (ga, gl)** at a given distance in pc

#### Parameters

- **gl** (*float, Angle, or Quantity*) – Galactic longitude in degrees (or astropy Angle)
- **gb** (*float, Angle, or Quantity*) – Galactic latitude in degrees (or astropy Angle)
- **dist** (*float or Quantity*) – Distance in pc

**Returns** electron density in cm<sup>-3</sup>

**Return type** N\_e (astropy.Quantity)

```
pygedm.ymw16_wrapper.calculate_electron_density_xyz(x, y, z)
```

Calculate electron density at a point with galactocentric coords (x, y, z)

#### Parameters

- **x** (*float or Quantity*) – galactocentric X coordinate in pc
- **y** (*float or Quantity*) – galactocentric Y coordinate in pc
- **z** (*float or Quantity*) – galactocentric Z coordinate in pc

**Returns** electron density in cm<sup>-3</sup>

**Return type** N\_e (astropy.quantity)

```
pygedm.ymw16_wrapper.dist_to_dm(gl, gb, dist, mode='gal', nu=1.0)
```

Convert a distance to a DM

#### Parameters

- **gl** (*float in deg or astropy.Angle*) – galactic longitude
- **gb** (*float in deg or astropy.Angle*) – galactic latitude
- **dist** (*float or astropy.Quantity*) – distance to source (pc) or if in mode IGM use (Mpc)
- **mode** (*str*) – Gal, MC, or IGM (for YMW16 only)

- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)

**Returns** dispersion measure (pc/cm<sup>3</sup>) and scattering time scale (s)

**Return type** dm (astropy.Quantity), tau\_sc (astropy.Quantity)

```
pygedm.ymw16_wrapper.dm_to_dist(gl, gb, dm, dm_host=0, mode='gal', nu=1.0)
```

Convert a DM to a distance

#### Parameters

- **gl** (*float in deg or astropy.Angle*) – galactic longitude
- **gb** (*float in deg or astropy.Angle*) – galactic latitude
- **dm** (*float in pc/cm<sup>3</sup> or astropy.Quantity*) – dispersion measure (pc cm<sup>-3</sup>)
- **mode** (*str*) – Gal, MC, or IGM (for YMW16 only)
- **nu** (*float in GHz or astropy.Quantity*) – observing frequency (GHz)

**Returns** distance (pc) and scattering time scale (s)

**Return type** dist (astropy.Quantity), tau\_sc (astropy.Quantity)

## pygedm.yt2020

Python implementation of Yamasaki & Totani DM Halo model

## References

[1] Yamasaki S, Totani T (2020), *The Galactic Halo Contribution to the Dispersion Measure of Extragalactic Fast Radio Bursts* The Astrophysical Journal, Volume 888, Issue 2, id.105

## Notes

Adapted from S. Yamasaki's DM\_halo\_yt2020\_numerical.py command-line python code

```
pygedm.yt2020.calculate_halo_dm(l, b, component='both')
```

Compute halo DM

#### Parameters

- **l** (*float*) – Galactic longitude, in degrees (-180 to +180)
- **b** (*float*) – Galactic latitude, in degrees (-90 to 90)
- **component** (*str*) – Compute ‘spherical’ component of halo, ‘disk’ component, or ‘both’ components.

**Returns** Dispersion measure in [pc/cm<sup>3</sup>]

**Return type** DM (float)

```
pygedm.yt2020.calculate_halo_dm_analytic(l, b)
```

Calculate the DM contribution of the Galactic halo.

Uses an analytical formula for speed. Useful for all-sky mapping.

#### Parameters

- **l** (*float*) – Galactic longitude, in degrees (-180 to +180)

- **b** (*float*) – Galactic latitude, in degrees (-90 to 90)

`pygedm.yt2020.ne_disk(l, b, s)`

Compute electron density for spherical component for (l, b) at distance s

### Parameters

- **l** (*float*) – Galactic longitude, in radians (-pi to +pi)
- **b** (*float*) – Galactic latitude, in radians (-pi/2 to pi/2)
- **s** (*float*) – Distance (kpc)

**Returns** electron density in cm<sup>-3</sup>

**Return type** ne (float)

`pygedm.yt2020.ne_sphe(l, b, s)`

Compute electron density for spherical component for (l, b) at distance s

### Parameters

- **l** (*float*) – Galactic longitude, in radians (-pi to +pi)
- **b** (*float*) – Galactic latitude, in radians (-pi/2 to pi/2)
- **s** (*float*) – Distance (kpc)

**Returns** electron density in cm<sup>-3</sup>

**Return type** ne (float)

`pygedm.yt2020.s_max(l, b)`

Compute integration limit s\_max for given sky coordinates

### Parameters

- **l** (*float*) – Galactic longitude, in radians (-pi to +pi)
- **b** (*float*) – Galactic latitude, in radians (-pi/2 to pi/2)

**Returns** s\_max (float), maximum integration limit corresponding to r = r\_vir

---

## Python Module Index

---

### p

`pygedm.ne2001_wrapper`, 8

`pygedm.pygedm`, 5

`pygedm.ymw16_wrapper`, 10

`pygedm.yt2020`, 11



### C

calculate\_electron\_density\_lbr () (in module `pygedm.pygedm`), 6  
calculate\_electron\_density\_lbr () (in module `pygedm.ymw16_wrapper`), 10  
calculate\_electron\_density\_xyz () (in module `pygedm.ne2001_wrapper`), 9  
calculate\_electron\_density\_xyz () (in module `pygedm.pygedm`), 6  
calculate\_electron\_density\_xyz () (in module `pygedm.ymw16_wrapper`), 10  
calculate\_halo\_dm () (in module `pygedm.pygedm`), 6  
calculate\_halo\_dm () (in module `pygedm.yt2020`), 11  
calculate\_halo\_dm\_analytic () (in module `pygedm.yt2020`), 11  
convert\_lbr\_to\_xyz () (in module `pygedm.pygedm`), 6

### D

dist\_to\_dm () (in module `pygedm.ne2001_wrapper`), 9  
dist\_to\_dm () (in module `pygedm.pygedm`), 7  
dist\_to\_dm () (in module `pygedm.ymw16_wrapper`), 10  
dm\_to\_dist () (in module `pygedm.ne2001_wrapper`), 9  
dm\_to\_dist () (in module `pygedm.pygedm`), 7  
dm\_to\_dist () (in module `pygedm.ymw16_wrapper`), 11

### G

generate\_healpix\_dm\_map () (in module `pygedm.pygedm`), 8

### N

ne\_disk () (in module `pygedm.yt2020`), 12  
ne\_sphe () (in module `pygedm.yt2020`), 12

### P

`pygedm.ne2001_wrapper` (module), 8  
`pygedm.pygedm` (module), 5  
`pygedm.ymw16_wrapper` (module), 10  
`pygedm.yt2020` (module), 11

### R

`run_from_pkgdir()` (in module `pygedm.ne2001_wrapper`), 10

### S

`s_max()` (in module `pygedm.yt2020`), 12

### T

`TAUISS()` (in module `pygedm.ne2001_wrapper`), 8